

EVALUATION OF TACHIYA'S ALGEBRAIC INFINITE PRODUCTS INVOLVING FIBONACCI AND LUCAS NUMBERS

Jonathan Sondow

209 West 97th Street #6F, New York, NY 10025, USA
jsondow@alumni.princeton.edu

Abstract. In 2007, Tachiya gave necessary and sufficient conditions for the transcendence of certain infinite products involving Fibonacci numbers F_k and Lucas numbers L_k . In the present note, we explicitly evaluate two classes of his algebraic examples. Special cases are

$$\prod_{n=1}^{\infty} \left(1 + \frac{1}{F_{2^n+1}}\right) = \frac{3}{\varphi}, \quad \prod_{n=1}^{\infty} \left(1 + \frac{1}{L_{2^n+1}}\right) = 3 - \varphi,$$

where $\varphi = (1 + \sqrt{5})/2$ is the golden ratio.

Mathematics Subject Classification 2000: 11B39, 11Y60

Keywords: Fibonacci number, Lucas number, infinite product, algebraic number

INTRODUCTION

For $k \geq 0$, define the *Fibonacci numbers* F_k by $F_{k+2} = F_{k+1} + F_k$ with $F_0 = 0$ and $F_1 = 1$, and the *Lucas numbers* L_k by $L_{k+2} = L_{k+1} + L_k$ with $L_0 = 2$ and $L_1 = 1$.

In 2003, D. Duverney and K. Nishioka [1] used Mahler's method to give a transcendence criterion for general series. As applications, they established necessary and sufficient conditions for the transcendence of the numbers

$$\sum_{n=0}^{\infty} \frac{c_n}{F_{r^n} + d_n}, \quad \sum_{n=0}^{\infty} \frac{c_n}{L_{r^n} + d_n},$$

where $r \geq 2$ is an integer and $\{c_n\}_{n \geq 0}$ and $\{d_n\}_{n \geq 0}$ are suitable sequences of algebraic numbers.

By modifying their method, Y. Tachiya [2] in 2007 proved analogous results for infinite products. In particular, he gave necessary and sufficient conditions for the transcendence of the numbers

$$\prod_{n=0}^{\infty} \left(1 + \frac{c_n}{F_{ar^n+b}}\right), \quad \prod_{n=0}^{\infty} \left(1 + \frac{c_n}{L_{ar^n+b}}\right),$$

where $r \geq 2$, a , and b are integers, $ar^n + b \neq 0$ for $n \geq 0$, and $\{c_n\}_{n \geq 0}$ is a suitable sequence of algebraic numbers. In the special case where the sequence is constant, his Examples 1 and 2 can be stated as the following two theorems.

Theorem 1 (Tachiya). *Let $a \geq 1$, $b \geq 0$, $r \geq 2$, and c be integers. Then*

$$\prod_{n=0}^{\infty} \left(1 + \frac{c}{F_{ar^n+b}} \right)$$

is algebraic if and only if

- (i) $c = 0$, or
- (ii) $r = 2$ and $c = F_b$.

In particular,

$$\prod_{n=0}^{\infty} \left(1 + \frac{c}{F_{r^n}} \right)$$

is algebraic if and only if $c = 0$.

Theorem 2 (Tachiya). *Let $a \geq 1$, $b \geq 0$, $r \geq 2$, and c be integers. Then*

$$\prod_{n=0}^{\infty} \left(1 + \frac{c}{L_{ar^n+b}} \right)$$

is algebraic if and only if

- (i) $c = 0$, or
- (ii) $r = 2$ and $c = L_b$, or
- (iii) $r = 2, b = 0$, and there exists a root of unity ω such that $\omega^{2^n} + \omega^{-2^n} = c$ for all large n .

In particular, Tachiya [2, p. 185] obtained that, for any integers $c \neq 0$ and $r \geq 2$, the number

$$\prod_{\substack{n=1 \\ L_{r^n} \neq -c}}^{\infty} \left(1 + \frac{c}{L_{r^n}} \right)$$

is transcendental, except for the two algebraic cases

$$\prod_{n=1}^{\infty} \left(1 + \frac{-1}{L_{2^n}} \right) = \frac{\sqrt{5}}{4}, \quad \prod_{n=1}^{\infty} \left(1 + \frac{2}{L_{2^n}} \right) = \sqrt{5},$$

which follow from the case (iii) with $\omega = (1 \pm \sqrt{-3})/2$ and $\omega = \pm 1$, respectively. He says, “These examples of algebraic infinite products involving Lucas numbers seem to be new.”

In the present note, we explicitly evaluate the algebraic infinite products in case (ii) of Theorems 1 and 2. These formulas may also be new.

THE FORMULAS

Proposition 1. *Let F_k and L_k be the k th Fibonacci and Lucas numbers, and let a and b be positive integers. Then*

$$\prod_{n=1}^{\infty} \left(1 + \frac{F_b}{F_{2^n a + b}} \right) = \frac{1 - (-1)^b \varphi^{-2a-2b}}{1 - \varphi^{-2a}} \quad (1)$$

and

$$\prod_{n=1}^{\infty} \left(1 + \frac{L_b}{L_{2^n a + b}} \right) = \frac{1 + (-1)^b \varphi^{-2a-2b}}{1 - \varphi^{-2a}}, \quad (2)$$

where

$$\varphi = \frac{1 + \sqrt{5}}{2}$$

is the golden ratio. In particular,

$$\prod_{n=1}^{\infty} \left(1 + \frac{1}{F_{2^n + 1}} \right) = \frac{3}{\varphi} \quad (3)$$

and

$$\prod_{n=1}^{\infty} \left(1 + \frac{1}{L_{2^n + 1}} \right) = 3 - \varphi. \quad (4)$$

Proof. (Compare [2, p. 199].) From Binet's formula

$$F_k = \frac{\varphi^k - (-\varphi^{-1})^k}{\varphi + \varphi^{-1}}$$

we have

$$\begin{aligned} \prod_{n=1}^{\infty} \left(1 + \frac{F_b}{F_{2^n a + b}} \right) &= \prod_{n=1}^{\infty} \left(1 + \frac{\varphi^b - (-1)^b \varphi^{-b}}{\varphi^{2^n a + b} - (-1)^b \varphi^{-2^n a - b}} \right) \\ &= \prod_{n=1}^{\infty} \left(1 + \frac{\varphi^{-2^n a - b} (\varphi^b - (-1)^b \varphi^{-b})}{1 - (-1)^b \varphi^{-2^{n+1} a - 2b}} \right) \\ &= \prod_{n=1}^{\infty} \left(1 + \varphi^{-2^n a} \right) \frac{1 - (-1)^b \varphi^{-2^n a - 2b}}{1 - (-1)^b \varphi^{-2^{n+1} a - 2b}} \\ &= \frac{1 - (-1)^b \varphi^{-2a-2b}}{1 - \varphi^{-2a}}, \end{aligned}$$

by cancellation and the fact that

$$\prod_{n=1}^{\infty} (1 + x^{2^n}) = \frac{1}{1 - x^2} \quad (|x| < 1).$$

This proves the equality (1). It and the relation $\varphi^2 = \varphi + 1$ yield

$$\prod_{n=1}^{\infty} \left(1 + \frac{1}{F_{2^n+1}} \right) = \frac{1 + \varphi^{-4}}{1 - \varphi^{-2}} = \frac{\varphi^4 + 1}{\varphi^4 - \varphi^2} = \frac{3\varphi + 3}{\varphi^2 + \varphi} = \frac{3}{\varphi},$$

proving (3). Using the formula $L_k = \varphi^k + (-\varphi)^{-k}$, the proofs of (2) and (4) are similar and will be omitted. $\square$

References

- [1] D. Duverney and K. Nishioka, “An inductive method for proving the transcendence of certain series”, *Acta Arith.* 110 (4), 305–330 (2003).
- [2] Y. Tachiya, “Transcendence of certain infinite products”, *J. Number Theory* 125, 182–200 (2007).